



THE HUMANE SOCIETY
OF THE UNITED STATES

An HSUS Report: Food Safety and Cage Egg Production

Abstract

States have begun legislating against cage egg production and dozens of major U.S. food retailers, restaurant chains, and foodservice providers—as well as hundreds of U.S. universities—are switching to cage-free eggs. Extensive scientific evidence strongly suggests this trend will improve food safety. The editor-in-chief of the trade journal *Egg Industry* agrees that assertions to the contrary are “invalid...unconvincing, unsupported and easily refuted.”¹ All ten scientific studies published in the last five years comparing *Salmonella* contamination between caged and cage-free operations found that those confining hens in cages had higher rates of *Salmonella*, the leading cause of food poisoning related death in the United States. This has led prominent consumer advocacy organizations, such as the Center for Food Safety, to oppose the use of cages to confine egg-laying hens.

Introduction

How we treat animals can have serious public health implications. The AIDS virus, for example, has killed more than 25 million people,² and its emergence has been traced to the butchering of chimpanzees for their flesh.³ The emergence of SARS, the contagious respiratory disease that infected thousands worldwide, has been linked to live animal markets,⁴ and the introduction of monkeypox into the United States has been blamed on the exotic pet trade.⁵ In fact, many of humanity’s great disease scourges—including smallpox,⁶ influenza,⁷ and measles⁸—likely originally arose from our domestication of farm animals.⁹

Many current industrial farming practices threaten the health of Americans, including the feeding of millions of pounds of antibiotics to farm animals every year.¹⁰ Antibiotics are routinely fed to farm animals in part to counteract stressful, overcrowded, and contaminated conditions found on factory farms.¹¹ Lacing the feed of farm animals with antibiotics has been condemned by the American Medical Association, the American Public Health Association, the Infectious Diseases Society of America, and the American Academy of Pediatrics, among 300 other organizations nationwide.¹² Despite the widespread outcry against this practice from the public health community, agribusiness continues to engage in this dangerous practice.

Other hazardous practices include the cannibalistic feeding of slaughterhouse waste, blood, and manure to farmed animals, blamed for the emergence of bovine spongiform encephalopathy (“mad cow disease”).¹³ The subsequent slaughter for human consumption of “downer” cows too sick or crippled to walk led to the largest meat recall in this country’s history.¹⁴

The intensive confinement of farm animals can also have negative public health implications.¹⁵ High stocking densities—the number of animals confined in a given space—have been associated with an elevated risk of infecting farm animals with a number of parasites and pathogens that can affect humans:

- *Yersinia enterocolitica* in goats;¹⁶
- *Trichostrongylus* in sheep;¹⁷
- *Mycobacterium bovis*,¹⁸ *Brucella*,¹⁹ *Salmonella*,²⁰ *Neospora*,²¹ and *Cryptosporidium* in cattle;²²
- *E. coli* O157:H7 in both sheep and cattle;²³
- *Ostertagia* in calves;²⁴

- *Oesophagostomum*,²⁵ Aujeszky's disease virus, and swine flu virus in pigs.²⁶

Farm animal well-being* and food safety issues are often inextricably linked.

Improvements in animal welfare can improve food safety by reducing stress-induced immunosuppression, infectious disease incidence, pathogen shedding, and antibiotic use and resistance.²⁷ Acute foodborne illnesses strike an estimated 76 million Americans every year²⁸ at an annual cost to the United States in excess of \$150 billion,²⁹ or nearly \$500 per American per year.³⁰ Studies show that small improvements in farm animal health may result in significant reductions in human illness.³¹

Eggborne *Salmonella*



Compassion Over Killing

Salmonella is the most commonly diagnosed foodborne bacterial illness in the United States³² and appears to be the leading cause of food-related death.³³ Eggs are the leading culprit for human *Salmonella* infection.³⁴ In 1994, a single egg-related outbreak sickened more than 200,000 Americans.³⁵ More typically, eggs may cause around 180,000 cases of *Salmonella* poisoning in the United States, according to scientists with the Centers for Disease Control and Prevention (CDC) and U.S. Department of Agriculture (USDA).³⁶ The annual cost of treating *Salmonella* infections in the billions.³⁷ As the U.S. Food and Drug Administration concluded in a 2009 press release: "Egg-associated illness caused by *Salmonella* is a serious public health problem."³⁸

Because *Salmonella* can infect the ovaries of hens, eggs from infected birds can be laid with the bacteria prepackaged inside.³⁹ According to research funded by the American Egg Board, *Salmonella* can survive sunny-side-up, over-easy, and scrambled cooking methods.⁴⁰ Infants and young children have been found to be at especially high risk.⁴¹ Although thousands die from food poisoning every year in the United States, the vast majority of victims suffer only acute, self-limited illnesses. *Salmonella*, however, can result in chronic arthritic joint inflammation⁴² and persistent irritable bowel syndrome in children.⁴³

Caged Hens Pose Significantly Higher *Salmonella* Risk

In U.S. commercial egg production, approximately 95% of laying hens are confined in battery cages, small wire enclosures that afford each hen roughly 430 cm² (67 in²)⁴⁴—a space smaller than a single sheet of letter-sized paper. These cages are placed side by side in rows and stacked in tiers commonly 4-8 levels high. Each cage may hold 5-10 birds⁴⁵ and hundreds of thousands of hens may be confined within a single building.

All 27 countries of the European Union (EU) are phasing out the use of these barren cages by 2012.⁴⁶ To study the public health implications of this move, an EU-wide *Salmonella* survey was launched in which more than 30,000 samples were taken from more than 5,000 operations across two dozen countries. This represents the best available data set comparing *Salmonella* infection risk between different laying hen housing systems. Without exception, for every *Salmonella* serotype grouping reported and for every type of production system examined, there was significantly higher *Salmonella* risk in operations that confine hens in cages.⁴⁷

* For information on the animal welfare implications, see "The Welfare of Intensively Confined Animals" at www.farmanimalwelfare.org



The European Food Safety Authority analysis found 43% lower odds of *Salmonella* Enteritidis contamination in cage-free barns, where hens are raised indoors, than in cage production. In organic egg production the odds of *Salmonella* contamination were 95% lower and in free range production the odds were 98% lower.⁵⁰ For *Salmonella* Typhimurium, the most common source of *Salmonella* poisoning in the United States,⁵¹ there was 77% lower odds of infection when hens were raised in barns compared to cages and 93% lower odds in organic and free range systems. For the other *Salmonella* serotypes found, compared to operations with hens in cages there was 96% lower odds in barn-raised flocks, 98% lower odds in

organic flocks, and 99% lower odds in free-ranging birds. This translates into at least 25-times greater odds of contamination on factory farms that confine hens in cages compared to cage-free production. The European Food Safety Authority analysis concluded: "Cage flock holdings are more likely to be contaminated with *Salmonella*."⁵²

Since this comprehensive survey was completed, ten additional studies have been published in the peer-reviewed scientific literature comparing *Salmonella* risk in caged and cage-free facilities. Without exception, each of them found higher rates of *Salmonella* in battery cage production units.^{53,54,55,56,57,58,59,60,61,62}

Every scientific study published in the last 5 years found increased <i>Salmonella</i> rates in cage operations
2010: 20 times greater odds of <i>Salmonella</i> infection in caged flocks
2010: 7.77 times greater odds of <i>Salmonella</i> Enteritidis in operations caging hens
2009: Significantly more risk of <i>Salmonella</i> in caged flocks
2008: 7.88 to 21.52 times greater odds of <i>Salmonella</i> in operations caging hens
2008: More than twice the prevalence of <i>Salmonella</i> in operations caging hens
2008: 59% greater prevalence of <i>Salmonella</i> (though not statistically significant)
2007: 1.8 to 25 times greater odds of <i>Salmonella</i> in operations caging hens
2007: 3.7 times greater prevalence of <i>Salmonella</i> in operations caging hens
2006: More than twice the prevalence of <i>Salmonella</i> in operations caging hens
2005: 1.9 to 6.7 times greater risk of <i>Salmonella</i> in typical ^{48,49} (dry manure) cages [†]

A recent article in the trade publication *World Poultry*, titled "*Salmonella* Thrives in Cage Housing," acknowledged that "the majority of the studies clearly indicate that a cage housing system has an increased risk of being *Salmonella*-positive in comparison to non-cage housing systems."⁶³ A study published in the December 2009 issue of *Poultry Science* found that cage-free hens experimentally infected with *Salmonella* may even clear the infection faster than caged hens.⁶⁴

The leading U.S. egg industry trade group claims that caging hens is "better for food safety,"⁶⁵ but in response to a landslide vote in California to ban the practice, the editor-in-chief of the trade journal *Egg Industry* admitted that such claims are "invalid...unconvincing, unsupportable and easily refuted."⁶⁶

Cage Production Factors That Increase *Salmonella* Risk

The reason cage operations have consistently been found to be at such higher risk for *Salmonella* is multifactorial. From the European Food Safety Authority analysis:

[†] Assuming a cage-free flock size of [20,000](#) versus a battery cage flock size of [100,000](#)

“In general, the higher prevalence [of *Salmonella*] in cage flocks might partly be explained by the fact that hens in the more intensive systems have a higher risk of being infected due to a relatively large flock size and higher density of hens. Moreover, cages can be difficult to disinfect and the housing may harbour breeding populations of rodents and other potential vectors such as flies or litter beetles. *Salmonella* has been shown to be more persistent in consecutive cage flocks compared with non-cage flocks in which the infection is more easily cleaned out during the empty period between flocks.”⁶⁷

Factor 1: Greater volume of fecal dust

Cage production facilities confine greater numbers of birds in a single building, as the caged birds are stacked in vertical tiers. There are single cage egg factories in the United States that cage millions of hens.⁶⁸ Such high densities of birds can produce a larger volume of contaminated airborne fecal dust, which may be responsible in part for the elevated threats to food safety posed by battery cage operations.⁶⁹ The latest national USDA survey of the domestic egg industry found that sheds confining more than 100,000 birds were four times more likely to be contaminated with *Salmonella*. The average number of hens confined in sheds found positive for *Salmonella* in the United States was 109,777,⁷⁰ much higher than cage-free operations typically hold.

Factor 2: More rodent disease vectors

The preponderance of disease-carrying rodents, flies, and other pests in battery cage sheds is another factor contributing to increased *Salmonella* infection rates in cage systems. Rodent infestations are closely tied to *Salmonella* rates.⁷¹ The manure pits typical of many cage operations are considered “ideal nesting grounds for rodents.”⁷² Indeed, rodents have been found to be “particularly persistent” in cage operations because they can breed in manure pits and “gain access to feeders, without interference from the birds,” who are confined in cages.⁷³ Other structural aspects of battery cage operations may also play a role in facilitating the rodent-borne spread of infection.⁷⁴

Factor 3: More insect disease vectors

According to the latest edition of Commercial Chicken Meat and Egg Production, the leading poultry science text,⁷⁵ one of many disadvantages of battery cage systems is that flies “are generally a greater nuisance” compared to cage-free production.⁷⁶ More than merely an annoyance, flies are proven vectors for *Salmonella* on egg farms.⁷⁷ According to Richard Axtell, a Professor Emeritus of Entomology: “By far the greatest populations of flies occur in the caged-layer houses that are widely used for commercial egg production.”⁷⁸ Scientists with the Food and Drug Administration agree: “In the poultry industry, the greatest numbers of houseflies and other disease-carrying flies occur in caged-layer houses (poultry houses with laying hens in cages for commercial egg production), where the flies breed in accumulated manure beneath the cages.”⁷⁹ In contrast, in cage-free broiler chicken houses, flies are “rarely a problem.”⁸⁰

Factor 4: Most difficult to disinfect

Salmonella can survive for more than two years in dried chicken feces,⁸¹ but can often be eliminated from laying hen houses with thorough cleaning and disinfection. Experts have noted, however, that cage operations are the “most difficult to clean properly”⁸² because of the “difficulty to efficiently disinfect the cages.”⁸³ The manure pits common in cage systems, which may not even be cleared between flocks, pose additional hygiene challenges.⁸⁴ From a poultry science journal:

“[C]age houses are intrinsically difficult to clean and disinfect to a good standard. Cages are normally organised in 3-12 tier stacks with associated complicated structures including dropping boards/belts drinkers, automatic egg belts, and feeder systems...Residual feed in particular may facilitate the multiplication of *Salmonella* after washing. In many cases older houses have no drainage, and electrical systems may not be water-proof. Because of these limitations, some buildings have only been ‘dry-cleaned’, which is normally...not satisfactory to achieve elimination of *Salmonella*.”⁸⁵

This has been validated in other countries. The Danish Veterinary and Food Administration states: “Experience shows that battery cage systems are particularly difficult to clean and disinfect.”⁸⁶ This is borne out by research

performed by the British Veterinary Laboratories Agency, which indicated “that there are particular problems with the disinfection of cage layer farms. This may be due to the larger flocks of birds kept at higher densities, which result in a larger volume of contaminated faecal material and dust, and the difficult access for cleaning in and around the cages.”⁸⁷

In comparison, cleaning and disinfecting equipment in cage-free facilities has been found to be more than twice as effective in combating *Salmonella* than attempts to disinfect battery cage operation equipment.⁸⁸ Even saturating a battery cage operation with formaldehyde-spiked steam for 24 consecutive hours at more than 140 degrees Fahrenheit—considered the gold standard treatment⁸⁹ found to effectively sterilize cage-free houses for *Salmonella*—may not effectively disinfect battery cage sheds.⁹⁰ To combat the rise of food poisoning caused by *Salmonella*, CDC researchers have called for a “sanitary revolution in farm-animal production.”⁹¹

Factor 5: More gut colonization and shedding

Research published in *Poultry Science* suggests another reason that chickens raised on bedding, rather than in bare, wire cages, have lower risk. On bedding, chickens may acquire natural gut flora that competitively prevents *Salmonella* colonization.⁹² Natural microflora would normally be obtained by chicks from their mothers and the environment. In industrial systems, however, chicks are no longer raised by hens but by incubators, after which they are confined in barren wire cages, potentially delaying or preventing the development of the normal adult gut flora helpful in preventing *Salmonella* infection.⁹³ Faster declines in *Salmonella* shedding have also been noted in experimentally infected cage-free hens compared to those confined in barren cages.⁹⁴

Factor 6: More stress

Physiological stress may also play a role.⁹⁵ In general, “the bulk of the evidence suggests that chronic or prolonged stress generally inhibits the immune response to infection, thus potentially rendering animals more susceptible to infectious disease.”⁹⁶ Specifically, research has shown that stress hormones can increase *Salmonella* colonization and systemic spread in chickens.⁹⁷ The stress hormone noradrenaline can boost the growth rate of *Salmonella* bacteria by orders of magnitude;⁹⁸ at the same time stress-related corticosteroids can impair the immune system.⁹⁹ A USDA researcher recently concluded that “there is increasing evidence to demonstrate that stress can have a significant deleterious effect on food safety.”¹⁰⁰

Increased Flock Risk Directly Translates To Increased Food Safety Risk

Contemporary studies universally show increased *Salmonella* rates in dust and manure samples from cage operations, providing conclusive evidence that measures to eliminate cages will likely improve the safety of the food supply. USDA researchers have found that “[f]locks with high levels of manure contamination were 10 times as likely to produce contaminated eggs as were flocks with low levels,” concluding that flocks with the highest levels of contamination “appeared to pose the greatest public health threat.”¹⁰¹ A key finding of a joint World Health Organization and Food and Agriculture Organization of the United Nations *Salmonella* risk assessment was that “[r]educing flock prevalence results in a directly proportional reduction in human health risk. For example, reducing flock prevalence from 50% to 25% results in a halving of the mean probability of illness per serving [of eggs].”¹⁰²

Infected hens can lay infected eggs. As of 2010, six studies have been published comparing *Salmonella* contamination rates of the eggs themselves from barren cage production versus typical cage-free systems. Half of the studies—one in Australia,¹⁰³ one in Brazil¹⁰⁴ and one in Ireland¹⁰⁵—found no *Salmonella* in eggs from either system. The other three found higher infection rates in eggs from caged hens compared to barn-raised birds.^{106,107,108}

In 1994-1995, a study was conducted at a California egg farm with both cage and cage-free housing systems, including three battery cage sheds and three cage-free barns. The prevalence of *Salmonella* in pooled egg samples from caged hens was nearly three times that of eggs from the cage-free (barn-raised) hens.¹⁰⁹ Though

the farm's free-range eggs were found to have higher rates, this was attributed to exceptional circumstances in that a creek "entirely composed of sewage effluent" bordered the property.¹¹⁰ More recently, the U.K. Food Standards Agency tested eggs from grocery stores. While 9 out of the 2,376 egg samples from caged hens came up positive for *Salmonella*, none of the 785 cartons of cage-free eggs tested was contaminated.¹¹¹ Testing foreign eggs coming into the country, the scientists found 132 of 1,329 samples of eggs from caged birds to be tainted with *Salmonella*, but, once again, none of the tested eggs from cage-free facilities were found to be positive with the pathogen.¹¹²

Type of eggs used	Cases		Controls		Odds ratio†	95% CI‡
	No.	%	No.	%		
White table eggs from battery flocks	48	19.8	38	9.4	2.4	1.5, 3.8
Brown table eggs from battery flocks	47	19.7	46	11.3	1.9	1.2, 3.0
Eggs from deep litter or free-range production	125	51.0	195	47.8	1.1	0.8, 1.6
Eggs from organic production	36	14.6	107	26.3	0.5	0.3, 0.8
Eggs sold at barnyards	49	19.8	109	26.5	0.6	0.4, 0.9
Pasteurized egg products	4	1.6	16	3.9	0.5	0.1, 1.4

Eating eggs from caged birds has been directly tied to human illness. In a 2002 prospective case-control study published in the *American Journal of Epidemiology*, people who recently ate eggs from caged hens had about twice the odds of being sickened by *Salmonella* compared to people who did not eat eggs from hens kept in cages. Those eating cage-free eggs were not at significantly elevated risk.¹¹³

American Journal of Epidemiology 156(7):654-61

The Industrialization of Egg Production Led To the *Salmonella* Pandemic

According to Dr. Robert Tauxe, the deputy director of the CDC's Division of Foodborne, Bacterial and Mycotic Diseases, foodborne *Salmonella* infections "became important public health concerns in parallel with the modern intensification of animal rearing...in the 1950s and 1960s in North America,"¹¹⁴ which is when U.S. egg industry began embracing cage systems.¹¹⁵ In the 1940s, *Salmonella* only sickened a few hundred Americans every year.¹¹⁶ Before the industrial intensification of egg production, *Salmonella* Enteritidis was not even found in eggs in the United States.¹¹⁷ By the beginning of the 21st century, however, *Salmonella* Enteritidis-contaminated eggs were sickening an estimated 182,000 Americans annually.¹¹⁸

In its landmark report, Emerging Infections: Microbial Threats to Health in the United States, the National Academy of Sciences' Institute of Medicine states that "the introduction of feedlots and large-scale poultry rearing and processing facilities has been implicated in the increasing incidence of human pathogens, such as *Salmonella*, in domestic animals over the past 30 years."¹¹⁹ There are many industrial practices that have contributed to the emergence of the eggborne *Salmonella* threat. For example, the egg industry's eradication of *Salmonella* Gallinarum, a serotype that primarily affects birds but not humans, may have created the ecological niche necessary for the emergence of *Salmonella* Enteritidis, which poses little threat to birds (and hence industry profits)¹²⁰ but sickens more than 100,000 Americans every year.¹²¹

Another contributory factor may be overcrowding. From the *Journal of the American Veterinary Medical Association*: "If salmonellae are inadvertently introduced into a large confinement rearing-production unit, a catastrophic epizootic [animal disease epidemic] might occur due to certain inherent environmental and stress factors, e.g. ...[o]verpopulation or crowding..."¹²² Professor John Evans, a poultry specialist and former FDA senior microbiologist, correctly predicted decades ago that "*Salmonella* infection of animals will occur more frequently and affect more individual animals as concentration of confinement increases."¹²³ U.S. caged hens are currently confined 5-10 birds per cage¹²⁴ affording each animal less living space than single sheet of letter-sized paper¹²⁵ for virtually their entire 1-2 year lifespan.¹²⁶

Factory farming practices not only led to the emergence of the eggborne *Salmonella* threat, but also to its global proliferation. It has been recognized for nearly 40 years that the "adoption of intensive rearing systems in the poultry and livestock industries today may create environments which encourage rapid spread of salmonella...infections...."¹²⁷ According to the World Health Organization, "[t]he factors facilitating the spread

of salmonellosis are associated with the intensification of animal and poultry production....¹²⁸ Specifically, these factors include industry's selective breeding practices,¹²⁹ the feeding of slaughterhouse waste to hens,¹³⁰ and forced starvation molting,¹³¹ which collectively placed the corporate interests of agribusiness above the safety of consumers by facilitating the spread of *Salmonella*.

Just as the feeding of dead animals to live ones triggered the mad cow disease crisis, this same practice has also been implicated in the worldwide spread of *Salmonella*.¹³² Once egg production wanes, hens may be ground up and rendered into what is called "spent hen meal," and then fed to other hens.¹³³ Annually, the United States has produced the majority¹³⁴ of the estimated 10 million tons of animal protein concentrates (such as meat, blood, and bone meal) incorporated worldwide into farm animal feed.¹³⁵

More than half the feed samples for farmed birds that contain slaughterhouse waste were found contaminated with *Salmonella* in FDA tests,¹³⁶ and numerous human *Salmonella* outbreaks have been specifically tied to feeding farm animals contaminated meat and bone meal.^{137,138,139} The use of manure in farm animal feed may have also played a role in the spread of *Salmonella*.¹⁴⁰ CDC researchers have estimated that more than 1,000,000 cases of *Salmonella* poisoning in America can be directly tied to feed containing animal byproducts.¹⁴¹

Industry Response to the Eggborne *Salmonella* Epidemic

U.S. efforts to control *Salmonella* pale in comparison to those made in other countries that have virtually eliminated the problem.¹⁴² In fact, attempts to enact similar farm safety programs domestically have been eliminated at the urging of the egg industry itself.¹⁴³ The U.S. industry trade group United Egg Producers openly praises its own efforts to obstruct public health measures, even publishing in its *Washington Report* that it added language to the USDA inspection budget that effectively killed the *Salmonella* testing program.¹⁴⁴ According to Marion Nestle, former U.S. Department of Health and Human Services Senior Nutrition Policy Advisor: "Major food industries oppose pathogen-control measures by every means at their disposal."¹⁴⁵

Rather than working to ensure the safety of their own products, the intensive farm animal industries have often tried shifting that responsibility to their own consumers. The safe-handling label on eggs is a prime example. "There has been a subtle turning of this on to the consumer," wrote Steve Bjerklie, former editor of *Meat and Poultry*, "and it's morally reprehensible."¹⁴⁶ Patricia Griffin, Chief of the Enteric Diseases Epidemiological Branch at the CDC, responded famously to this blame-the-victim attitude with regard to *E. coli* O157:H7, another dangerous pathogen. "Is it reasonable," she asked, "that if a consumer undercooks a hamburger...their three-year-old dies?"¹⁴⁷ *Salmonella* has been estimated to kill 10 times more Americans every year than *E. coli* O157:H7.¹⁴⁸

Animal agribusiness understands that profitable but risky practices like these must be kept hidden from the public. "One of the best things modern animal agriculture has going for it is that most people...haven't a clue how animals are raised and processed," wrote an editor of the *Journal of Animal Science* in an animal agriculture textbook. "For modern animal agriculture, the less the consumer knows about what's happening before the meat hits the plate, the better."¹⁴⁹

Pew Commission on Industrial Farm Animal Production

The Pew Commission on Industrial Farm Animal Production was formed to conduct a comprehensive, fact-based, and balanced examination of key aspects of the farm animal industry. This prestigious independent panel was chaired by former Kansas Governor John Carlin and included former U.S. Secretary of Agriculture Dan Glickman, former Assistant Surgeon General Michael Blackwell, and James Merchant, then Dean of the University of Iowa College of Public Health, among many other experts across several disciplines. After a rigorous two-and-one-half-year inquiry, the Commissioners emphasized that the "ethical treatment of animals raised for food is essential to, and consistent with, achieving a safe and sustainable system for producing food

animals”¹⁵⁰ and concluded that “[d]ue to the large numbers of animals housed in close quarters in typical [industrial farm animal production] facilities there are many opportunities for animals to be infected by several strains of pathogens, leading to increased chance for a strain to emerge that can infect and spread in humans.”¹⁵¹

The Commissioners affirmed that “[f]ood animals that are treated well and provided with at least minimum accommodation of their natural behaviors and physical needs are healthier and safer for human consumption.”¹⁵² Specifically, they asserted that “[p]ractices that restrict natural motion...induce high levels of stress in the animals and threaten their health, which in turn may threaten human health.”¹⁵³ The Pew Commission on Industrial Farm Animal Production unanimously concluded that battery cages should be eliminated from American agriculture.¹⁵⁴

Conclusion

Institutions, corporations, electorates, and legislatures are increasingly embracing the recommendations of the Pew Commission. In a landslide 2008 vote, California moved to criminalize the cage confinement of egg-laying hens.¹⁵⁵ In 2009, Michigan's governor signed legislation that similarly phases out battery cages,¹⁵⁶ and in 2010 Governor Schwarzenegger signed a bill to ban the sale of shell eggs from caged hens imported into California.¹⁵⁷ Supermarket chains such as Safeway, Harris Teeter, Winn-Dixie, Trader Joe's, and Whole Foods have all taken steps to increase their sales of cage-free eggs. All of Wal-Mart's private label eggs are now cage-free. National restaurant chains, including Burger King, Wendy's, Denny's, Red Robin, Subway, Quiznos, Sonic, Hardee's and Carl's Jr., have also started using cage-free eggs. Compass Group, the world's largest food service provider switched to cage-free eggs for all of its 7,500 U.S. clients.

The best available science suggests that confining birds in cages means increased *Salmonella* infection risk in the birds, their eggs, and the consumers of caged eggs. The cage-free trend in the United States is therefore expected to increase the safety of the American food supply.

¹ Shane S. 2008. Proposition 2: Isolated anomaly...or national trend? Egg Industry, December, p. 4. www.eggindustry-digital.com/eggindustry/200812/#pg4. Accessed March 15, 2010.

² UNAIDS. 2006. Report on the global AIDS epidemic. www.unaids.org/en/KnowledgeCentre/HIVData/GlobalReport/2006/default.asp. Accessed March 15, 2010.

³ Sharp PM, Bailes E, Chaudhuri RR, Rodenburg CM, Santiago MO, and Hahn BH. 2001. The origins of acquired immune deficiency syndrome viruses: where and when? *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* 356(1410):867-76.

⁴ Kan B, Wang M, Jing H, et al. 2005. Molecular evolution analysis and geographic investigation of severe acute respiratory syndrome coronavirus-like virus in palm civets at an animal market and on farms. *Journal of Virology* 79(18):11892-900.

⁵ Gross E. 2003. Update: Multistate outbreak of monkeypox—Illinois, Indiana, Kansas, Missouri, Ohio, and Wisconsin, 2003. *Annals of Emergency Medicine* 42(5):660-4.

⁶ Gubser C, Hue S, Kellam P, and Smith GL. 2004. Poxvirus genomes: a phylogenetic analysis. *Journal of General Virology* 85(1):105-17.

⁷ Shortridge KF. 2003. Severe acute respiratory syndrome and influenza: virus incursions from southern China. *American Journal of Respiratory and Critical Care Medicine* 168(12):1416-20.

⁸ Weiss RA. 2001. The Leeuwenhoek Lecture 2001. Animal origins of human infectious disease. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* 356(1410):957-77.

⁹ Wolfe ND, Dunavan CP, and Diamond J. 2007. Origins of major human infectious diseases. *Nature* 447(7142):279-83.

¹⁰ Mellon M, Benbrook C, and Benbrook KL. 2001. Hogging It: Estimates of Antimicrobial Abuse in Livestock (Cambridge, MA: Union of Concerned Scientists). www.ucsusa.org/food_and_environment/antibiotics_and_food/hogging-it-estimates-of-antimicrobial-abuse-in-livestock.html. Accessed March 15, 2010.

¹¹ Office of Technology Assessment. 1979. *Drugs in Livestock Feed. Volume 1: Technical Report* (Washington, DC: U.S. Government Printing Office). http://govinfo.library.unt.edu/ota/Ota_5/DATA/1979/7905.PDF. Accessed March 15, 2010.

-
- ¹² Keep Antibiotics Working. 2007. Kennedy, Snowe & Slaughter introduce AMA-backed bill to cut antibiotic resistance linked to misuse of antibiotics in animal agriculture. Press release issued February 12. www.KeepAntibioticsWorking.com/new/resources_library.cfm?ReffID=97314. Accessed March 15, 2010.
- ¹³ Wilesmith JW, Ryan JB, and Atkinson MJ. 1991. Bovine spongiform encephalopathy: epidemiological studies on the origin. *The Veterinary Record* 128(9):199-203.
- ¹⁴ U.S. Department of Agriculture Food Safety and Inspection Service. 2008. California firm recalls beef products derived from non-ambulatory cattle without the benefit of proper inspection. Press release issued February 17. www.fsis.usda.gov/pdf/recall_005-2008_Release.pdf. Accessed March 15, 2010.
- ¹⁵ Tauxe RV. 2002. Emerging foodborne pathogens. *International Journal of Food Microbiology* 78(1-2):31-41.
- ¹⁶ Lanada EB, Morris RS, Jackson R, and Fenwick SG. 2005. Prevalence of *Yersinia* species in goat flocks. *Australian Veterinary Journal* 83(9):563-6.
- ¹⁷ Thamsborg SM, Jørgensen RJ, Waller PJ, and Nansen P. 1996. The influence of stocking rate on gastrointestinal nematode infections of sheep over a 2-year grazing period. *Veterinary Parasitology* 67(3-4):207-24.
- ¹⁸ White PC and Benhin JK. 2004. Factors influencing the incidence and scale of bovine tuberculosis in cattle in southwest England. *Preventive Veterinary Medicine* 63(1-2):1-7.
- ¹⁹ Salman MD, and Meyer ME. 1984. Epidemiology of bovine brucellosis in the Mexicali Valley, Mexico: Literature review of disease-associated factors. *American Journal of Veterinary Research* 45(8): 1557-1560.
- ²⁰ Jones PW, Collins P, Brown GT, and Aitken MM. 1983. *Salmonella* Saint-Paul infection in two dairy herds. *The Journal of Hygiene* 91(2):243-57.
- ²¹ Sanderson MW, Gay JM, and Baszler TV. 2000. *Neospora caninum* seroprevalence and associated risk factors in beef cattle in the northwestern United States. *Veterinary Parasitology* 90(1-2):15-24.
- ²² Atwill ER, Johnson EM, and Pereira MG. 1999. Association of herd composition, stocking rate, and duration of calving season with fecal shedding of *Cryptosporidium parvum* oocysts in beef herds. *Journal of the American Veterinary Medical Association* 215(12):1833-8.
- ²³ Stacey KF, Parsons DJ, Christiansen KH, and Burton CH. 2007. Assessing the effect of interventions on the risk of cattle and sheep carrying *Escherichia coli* O157:H7 to the abattoir using a stochastic model. *Preventive Veterinary Medicine* 79(1):32-45.
- ²⁴ Nansen P, Foldager J, Hansen JW, Henriksen SA, and Jørgensen RJ. 1988. Grazing pressure and acquisition of *Ostertagia ostertagi* in calves. *Veterinary Parasitology* 27(3-4):325-35.
- ²⁵ Thamsborg SM, Roepstorff A, and Larsen M. 1999. Integrated and biological control of parasites in organic and conventional production systems. *Veterinary Parasitology* 84(3-4):169-86.
- ²⁶ Maes D, Deluyker H, Verdonck M, et al. 2000. Herd factors associated with the seroprevalences of four major respiratory pathogens in slaughter pigs from farrow-to-finish pig herds. *Veterinary Research* 31(3):313-27.
- ²⁷ de Passillé AM and Rushen J. Food safety and environmental issues in animal welfare. *Revue Scientifique et Technique de l'Office International des Epizooties* 24(2):757-66.
- ²⁸ Mead PS, Slutsker L, Dietz V, et al. 1999. Food-related illness and death in the United States. *Emerging Infectious Diseases* 5(5):607-25.
- ²⁹ Scharff RL. 2010. Health-Related Costs from Foodborne Illness in the United States. March 3. www.producesafetyproject.org/admin/assets/files/Health-Related-Foodborne-Illness-Costs-Report.pdf-1.pdf. Accessed 2 August 2010.
- ³⁰ U.S. Census Bureau. 2010. Population clocks. www.census.gov/. Accessed 2 August 2010.
- ³¹ Singer RS, Cox LA Jr, Dickson JS, et al. Modeling the relationship between food animal health and human foodborne illness. *Preventive Veterinary Medicine* 79(2-4):186-203.
- ³² Chittick P, Sulka A, Tauxe RV, and Fry AM. 2006. A summary of national reports of foodborne outbreaks of *Salmonella* Heidelberg infections in the United States: clues for disease prevention. *Journal of Food Protection* 69(5):1150-3.
- ³³ Mead PS, Slutsker L, Dietz V, et al. 1999. Food-related illness and death in the United States. *Emerging Infectious Diseases* 5(5):607-25.
- ³⁴ Patrick ME, Adcock PM, Gomez TM, et al. 2004. *Salmonella* Enteritidis infections, United States, 1985-1999. *Emerging Infectious Diseases* 10(1):1-7.
- ³⁵ Hennessy TW, Hedberg CW, Slutsker L, et al. 1996. A national outbreak of *Salmonella* Enteritidis infections from ice cream. *The New England Journal of Medicine* 334(20):1281-6.
- ³⁶ Schroeder CM, Naugle AL, Schlosser WD, et al. 2005. Estimate of illnesses from *Salmonella* Enteritidis in eggs, United States, 2000. *Emerging Infectious Diseases* 11(1):113-5.
- ³⁷ Bryan FL and Doyle MP. 1995. Health risks and consequences of *Salmonella* and *Campylobacter jejuni* in raw poultry. *Journal of Food Protection* 58(3):326-44.

- ³⁸ U.S. Food and Drug Administration. 2009. New public health regulation to improve egg safety and reduce salmonella illnesses. Press release issued July 7. www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm170788.htm. Accessed March 15, 2010.
- ³⁹ Gast RK and Beard CW. 1990. Production of *Salmonella* Enteritidis-contaminated eggs by experimentally infected hens. *Avian Diseases* 34(2):438-46.
- ⁴⁰ Davis AL, Curtis PA, Conner DE, McKee SR, and Kerth LK. 2008. Validation of cooking methods using shell eggs inoculated with *Salmonella* serotypes Enteritidis and Heidelberg. *Poultry Science* 87(8):1637-42.
- ⁴¹ Trevejo RT, Courtney JG, Starr M, and Vugia DJ. 2003. Epidemiology of salmonellosis in California, 1990-1999: morbidity, mortality, and hospitalization costs. *American Journal of Epidemiology* 157(1):48-57.
- ⁴² Ternhag A, Törner A, Svensson A, Ekdahl K, and Giesecke J. 2008. Short- and long-term effects of bacterial gastrointestinal infections. *Emerging Infectious Diseases* 14(1):143-8.
- ⁴³ Saps M, Pensabene L, Di Martino L, et al. 2008. Post-infectious functional gastrointestinal disorders in children. *The Journal of Pediatrics* 152(6):812-6.
- ⁴⁴ United Egg Producers. 2010. United Egg Producers Animal Husbandry Guidelines for U.S. Egg Laying Flocks, 2010 Edition (Alpharetta, GA: United Egg Producers). www.uepcertified.com/media/pdf/UEP-Animal-Welfare-Guidelines.pdf. Accessed January 23, 2010.
- ⁴⁵ Bell DD and Weaver WD. 2002. *Commercial Chicken Meat and Egg Production*, 5th Edition (Norwell, MA: Kluwer Academic Publishers, p.1009).
- ⁴⁶ European Commission. 1999. Animal welfare on the farm—laying hens. http://ec.europa.eu/food/animal/welfare/farm/laying_hens_en.htm. Accessed December 30, 2009.
- ⁴⁷ European Food Safety Authority. 2007. Report of the Task Force on Zoonoses Data Collection on the Analysis of the baseline study on the prevalence of *Salmonella* in holdings of laying hen flocks of *Gallus gallus*. The EFSA Journal 97. www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178620761896.htm. Accessed March 15, 2010.
- ⁴⁸ U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Veterinary Services. 2000. Part II: Reference of 1999 Table Egg Layer Management in the U.S.. Layers '99, p. 42. nahms.aphis.usda.gov/poultry/layers99/Layers99_dr_PartII.pdf. Accessed August 24, 2010.
- ⁴⁹ Spelling FR and Whiting NE. 2007. *Environmental Management of Concentrated Animal Feeding Operations (CAFOs)* (Boca Raton, FL: CRC Press, p. 387).
- ⁵⁰ European Food Safety Authority. 2007. Report of the Task Force on Zoonoses Data Collection on the Analysis of the baseline study on the prevalence of *Salmonella* in holdings of laying hen flocks of *Gallus gallus*. The EFSA Journal 97. www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178620761896.htm. Accessed March 15, 2010.
- ⁵¹ Centers for Disease Control and Prevention. 2006. Preliminary FoodNet data on the incidence of infection with pathogens transmitted commonly through food--10 States, United States, 2005. *Morbidity and Mortality Weekly Report* 55(14):392-5.
- ⁵² European Food Safety Authority. 2007. Report of the Task Force on Zoonoses Data Collection on the Analysis of the baseline study on the prevalence of *Salmonella* in holdings of laying hen flocks of *Gallus gallus*. The EFSA Journal 97. www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178620761896.htm. Accessed March 15, 2010.
- ⁵³ Van Hoorebeke S, Van Immerseel F, Schulz J, et al. 2010. Determination of the within and between flock prevalence and identification of risk factors for *Salmonella* infections in laying hen flocks housed in conventional and alternative systems. *Preventive Veterinary Medicine* 94(1-2):94-100.
- ⁵⁴ Snow LC, Davies RH, Christiansen KH, et al. 2010. Investigation of risk factors for *Salmonella* on commercial egg-laying farms in Great Britain, 2004-2005. *Veterinary Record* 166(19):579-86.
- ⁵⁵ Huneau-Salaün A, Chemaly M, Le Bouquin S, et al. 2009. Risk factors for *Salmonella* enterica subsp. Enteric contamination in 5 French laying hen flocks at the end of the laying period. *Preventive Veterinary Medicine* 89:51-8.
- ⁵⁶ Namata H, Méroc E, Aerts M, et al. 2008. *Salmonella* in Belgian laying hens: an identification of risk factors. *Preventive Veterinary Medicine* 83(3-4):323-36.
- ⁵⁷ Mahé A, Bougeard S, Huneau-Salaün A, et al. 2008. Bayesian estimation of flock-level sensitivity of detection of *Salmonella* spp., Enteritidis and Typhimurium according to the sampling procedure in French laying-hen houses. *Preventive Veterinary Medicine* 84(1-2):11-26.
- ⁵⁸ Pieskus J, et al. 2008. *Salmonella* incidence in broiler and laying hens with the different housing systems. *Journal of Poultry Science* 45:227-231.
- ⁵⁹ European Food Safety Authority. 2007. Report of the Task Force on Zoonoses Data Collection on the Analysis of the baseline study on the prevalence of *Salmonella* in holdings of laying hen flocks of *Gallus gallus*. The EFSA Journal 97. www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178620761896.htm. Accessed March 15, 2010.
- ⁶⁰ Snow LC, Davies RH, Christiansen KH, et al. 2007. Survey of the prevalence of *Salmonella* species on commercial laying farms in the United Kingdom. *The Veterinary Record* 161(14):471-6.

-
- ⁶¹ Methner U, Diller R, Reiche R, and Böhland K. 2006. [Occurrence of salmonellae in laying hens in different housing systems and inferences for control]. *Berliner und Münchener tierärztliche Wochenschrift* 119(11-12):467-73.
- ⁶² Mollenhorst H, van Woudenberg CJ, Bokkers EG, de Boer IJ. 2005. Risk factors for *Salmonella enteritidis* infections in laying hens. *Poultry Science* 84(8):1308-13.
- ⁶³ 2009. *Salmonella* thrives in cage housing. *World Poultry* 25(10):18-9.
- ⁶⁴ De Vylder J, Van Hoorebeke S, Ducatelle R, et al. 2009. Effect of the housing system on shedding and colonization of gut and internal organs of laying hens with *Salmonella* Enteritidis. *Poultry Science* 88:2491-5
- ⁶⁵ Gregory C. 2009. Letter to members of United Egg Producers. www.unitedegg.org/. Accessed March 15, 2010.
- ⁶⁶ Shane S. 2008. Proposition 2: Isolated anomaly...or national trend?. *Egg Industry*, December, p. 4. www.eggindustry-digital.com/eggindustry/200812/#pg4. Accessed March 15, 2010.
- ⁶⁷ European Food Safety Authority. 2007. Report of the Task Force on Zoonoses Data Collection on the Analysis of the baseline study on the prevalence of *Salmonella* in holdings of laying hen flocks of *Gallus gallus*. The EFSA Journal 97. www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178620761896.htm. Accessed March 15, 2010.
- ⁶⁸ Ohio Department of Agriculture Livestock Environmental Permitting Program. 2010. www.agri.ohio.gov/apps/lepp_permits/lepp_permits.aspx. Accessed April 9, 2010.
- ⁶⁹ Namata H, Méroc E, Aerts M, et al. 2008. *Salmonella* in Belgian laying hens: an identification of risk factors. *Preventive Veterinary Medicine* 83(3-4):323-36.
- ⁷⁰ U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Veterinary Services. 2000. *Salmonella enterica* serotype Enteritidis in table egg layers in the U.S. National Animal Health Monitoring System, Layers '99. http://nahms.aphis.usda.gov/poultry/layers99/Layers99_dr_Salmonella.pdf. Accessed March 15, 2010.
- ⁷¹ Garber L, Smeltzer M, Fedorka-Cray P, Ladely S, and Ferris K. 2003. *Salmonella enterica* serotype Enteritidis in table egg layer house environments and in mice in U.S. layer houses and associated risk factors. *Avian Diseases* 47(1):134-42.
- ⁷² Carrique-Mas JJ and Davies RH. 2008. *Salmonella* Enteritidis in commercial layer flocks in Europe: legislative background, on-farm sampling and main challenges. *Brazilian Journal of Poultry Science* 10(1):1-9.
- ⁷³ Davies RH. 2005. Pathogen populations on poultry farms. In: Mead GC (ed.), *Food Safety Control in the Poultry Industry* (Cambridge, England: Woodhead Publishing Limited, p. 114).
- ⁷⁴ Carrique-Mas JJ and Davies RH. 2008. *Salmonella* Enteritidis in commercial layer flocks in Europe: legislative background, on-farm sampling and main challenges. *Brazilian Journal of Poultry Science* 10(1):1-9.
- ⁷⁵ Dale N. 2002. Book review: *Commercial Chicken Meat and Egg Production*. *The Journal of Applied Poultry Research* 11(2):224-5.
- ⁷⁶ Bell DD. 2001. Cage management for layers. In: Bell DD and Weaver WD Jr (eds.), *Commercial Chicken Meat and Egg Production*, 5th Edition (Norwell, MA: Kluwer Academic Publishers).
- ⁷⁷ Olsen AR and Hammack TS. 2000. Isolation of *Salmonella* spp. from the housefly, *Musca domestica* L., and the dump fly, *Hydrotaea aenescens* (Wiedemann) (Diptera: Muscidae), at caged-layer houses. *Journal of Food Protection* 63(7):958-60.
- ⁷⁸ Axtell RC and Arends JJ. 1990. Ecology and management of arthropod pests of poultry. *Annual Review of Entomology* 35:101-26.
- ⁷⁹ Olsen AR and Hammack TS. 2000. Isolation of *Salmonella* spp. from the housefly, *Musca domestica* L., and the dump fly, *Hydrotaea aenescens* (Wiedemann) (Diptera: Muscidae), at caged-layer houses. *Journal of Food Protection* 63(7):958-60.
- ⁸⁰ Axtell RC and Arends JJ. 1990. Ecology and management of arthropod pests of poultry. *Annual Review of Entomology* 35:101-26.
- ⁸¹ Davies RH and Breslin M. 2003. Persistence of *Salmonella* Enteritidis Phage Type 4 in the environment and arthropod vectors on an empty free-range chicken farm. *Environmental Microbiology* 5(2):79-84.
- ⁸² Gradel KO. 2004. Disinfection of *Salmonella* in poultry houses. Ph.D. thesis, February. University of Bristol Department of Clinical Veterinary Science.
- ⁸³ Namata H, Méroc E, Aerts M, et al. 2008. *Salmonella* in Belgian laying hens: an identification of risk factors. *Preventive Veterinary Medicine* 83(3-4):323-36.
- ⁸⁴ Carrique-Mas JJ and Davies RH. 2008. *Salmonella* Enteritidis in commercial layer flocks in Europe: legislative background, on-farm sampling and main challenges. *Brazilian Journal of Poultry Science* 10(1):1-9.
- ⁸⁵ Carrique-Mas JJ and Davies RH. 2008. *Salmonella* Enteritidis in commercial layer flocks in Europe: legislative background, on-farm sampling and main challenges. *Brazilian Journal of Poultry Science* 10(1):1-9.
- ⁸⁶ The Danish Veterinary and Food Administration. 2004. The national *Salmonella* control programme for the production of table eggs and broilers 1996-2002. *Fødevare Rapport* 6, March.
- ⁸⁷ Davies R and Breslin M. 2003. Observations on *Salmonella* contamination of commercial laying farms before and after cleaning and disinfection. *The Veterinary Record* 152(10):283-7.

-
- ⁸⁸ Davies R and Breslin M. 2003. Observations on *Salmonella* contamination of commercial laying farms before and after cleaning and disinfection. *The Veterinary Record* 152(10):283-7.
- ⁸⁹ Gradel KO. 2004. Disinfection of *Salmonella* in poultry houses. Ph.D. thesis, February. University of Bristol Department of Clinical Veterinary Science.
- ⁹⁰ Gradel KO, Jørgensen JC, Andersen JS, and Corry JEL. 2004. Monitoring the efficacy of steam and formaldehyde treatment of naturally *Salmonella*-infected layer houses. *Journal of Applied Microbiology* 96(3):613-22.
- ⁹¹ Crump JA, Griffin PM, and Angulo FJ. 2002. Bacterial contamination of animal feed and its relationship to human foodborne illness. *Clinical Infectious Diseases* 35(7):859-65.
- ⁹² Santos FB, Sheldon BW, Santos AA Jr, and Ferket PR. 2008. Influence of housing system, grain type, and particle size on *Salmonella* colonization and shedding of broilers fed triticale or corn-soybean meal diets. *Poultry Science* 87(3):405-20.
- ⁹³ Reynolds D. 2004. Tenants of the last 1.5 metres. *Microbiologist* 5(3):26-30.
- ⁹⁴ De Vylder J, Van Hoorebeke S, Ducatelle R, et al. 2009. Effect of the housing system on shedding and colonization of gut and internal organs of laying hens with *Salmonella* Enteritidis. *Poultry Science* 88:2491-5
- ⁹⁵ Humphrey T. 2006. Are happy chickens safer chickens? Poultry welfare and disease susceptibility. *British Poultry Science* 47(4):379-91.
- ⁹⁶ de Passillé AM and Rushen J. Food safety and environmental issues in animal welfare. *Revue Scientifique et Technique de l'Office International des Epizooties* 24(2):757-66.
- ⁹⁷ Methner U, Rabsch W, Reissbrodt R, and Williams PH. 2008. Effect of norepinephrine on colonisation and systemic spread of *Salmonella enterica* in infected animals: Role of catechol siderophore precursors and degradation products. *International Journal of Medical Microbiology* 298(5-6):429-39.
- ⁹⁸ Bailey MT, Karaszewski JW, Lubach GR, Coe CL, and Lyte M. 1999. In vivo adaptation of attenuated *Salmonella* Typhimurium results in increased growth upon exposure to norepinephrine. *Physiology and Behavior* 67(3):359-64.
- ⁹⁹ Shini S, Kaiser P, Shini A, and Bryden WL. 2008. Biological response of chickens (*Gallus gallus domesticus*) induced by corticosterone and a bacterial endotoxin. *Comparative Biochemistry and Physiology. Part B, Biochemistry & Molecular Biology* 149(2):324-33.
- ¹⁰⁰ Rostagno MH. 2009. Can stress in farm animals increase food safety risk? *Foodborne Pathogens and Disease* 6(7):767-76.
- ¹⁰¹ Henzler DJ, Kradel DC, and Sicho WM. 1998. Management and environmental risk factors for *Salmonella enteritidis* contamination of eggs. *American Journal of Veterinary Research* 59(7):824-9.
- ¹⁰² World Health Organization and the Food and Agriculture Organization of the United Nations. 2002. Risk assessments of *Salmonella* in eggs and broiler chickens. Microbiological risk assessment series 2. www.fao.org/DOCREP/005/Y4392E/Y4392E00.HTM. Accessed March 15, 2010.
- ¹⁰³ Barnett JL. 1998. The welfare and productivity of hens in a barn system and cages. A report for the Rural Industries Research and Development Corporation.
- ¹⁰⁴ Barbosa Filho JAD, Silva MAN, Silva IJO, and Coelho AAD. 2005. Egg quality in layers housed in different production systems and submitted to two environmental conditions. *Brazilian Journal of Poultry Science* 8(1):23-8.
- ¹⁰⁵ Food Safety Authority of Ireland. 2003. Bacteriological safety of eggs produced under the Bord Bia Egg Quality Assurance Scheme (EQAS).
- ¹⁰⁶ Kinde H, Read DH, Chin RP, et al. 1996. *Salmonella* Enteritidis, phage type 4 infection in a commercial layer flock in southern California: bacteriologic and epidemiologic findings. *Avian Diseases* 40(3):665-71.
- ¹⁰⁷ U.K. Food Standards Agency. 2004. Report of the survey of *Salmonella* contamination of U.K. produced shell eggs on retail sale. March 18. www.food.gov.uk/multimedia/pdfs/fsis5004report.pdf. Accessed March 15, 2010.
- ¹⁰⁸ Little CL, Walsh S, Hucklesby L, et al. 2006. Survey of *Salmonella* contamination of non-U.K. produced shell eggs on retail sale in the north west of England and London. Final report - Project B18012, November 15. U.K. Food Standards Agency.
- ¹⁰⁹ Kinde H, Read DH, Chin RP, et al. 1996. *Salmonella* Enteritidis, phage type 4 infection in a commercial layer flock in southern California: bacteriologic and epidemiologic findings. *Avian Diseases* 40(3):665-71.
- ¹¹⁰ Kinde H, Read DH, Ardans A, et al. 1996. Sewage effluent: likely source of *Salmonella* Enteritidis, phage type 4 infection in a commercial chicken layer flock in southern California. *Avian Diseases* 40(3):672-6.
- ¹¹¹ U.K. Food Standards Agency. 2004. Report of the survey of *Salmonella* contamination of U.K. produced shell eggs on retail sale. March 18. www.food.gov.uk/multimedia/pdfs/fsis5004report.pdf. Accessed March 15, 2010.
- ¹¹² Little CL, Walsh S, Hucklesby L, et al. 2006. Survey of *Salmonella* contamination of non-U.K. produced shell eggs on retail sale in the north west of England and London. Final report - Project B18012, November 15. U.K. Food Standards Agency.

-
- ¹¹³ Mølbak K and Neimann J. 2002. Risk factors for sporadic infection with *Salmonella* Enteritidis, Denmark, 1997-1999. *American Journal of Epidemiology* 156(7):654-61.
- ¹¹⁴ Tauxe RV. 1999. *Salmonella* Enteritidis: the continuing global public health challenge. In: Saeed AM, Gast RK, Potter ME, and Wall PG (eds.), *Salmonella enterica* Serovar Enteritidis in Humans and Animals: Epidemiology, Pathogenesis, and Control (Ames, IA: Iowa State University Press, pp. xi-xiii).
- ¹¹⁵ Van de Poel I. 1998. Why are chickens housed in battery cages? In: Disco C and van der Meulen B (eds.), *Studies in Making Sociotechnical Order* (New York, NY: Walter de Gruyter, pp. 143-177).
- ¹¹⁶ Morse EV and Duncan MA. 1974. Salmonellosis—an environmental health problem. *Journal of the American Veterinary Medical Association* 165(11):1015-9.
- ¹¹⁷ Rabsch W, Tschäpe H, and Bäumlner AJ. 2001. Non-typhoidal salmonellosis: emerging problems. *Microbes and Infection* 3(3):237-47.
- ¹¹⁸ Schroeder CM, Naugle AL, Schlosser WD, et al. 2005. Estimate of illnesses from *Salmonella* Enteritidis in eggs, United States, 2000. *Emerging Infectious Diseases* 11(1):113-5.
- ¹¹⁹ Lederberg J, Shope RE, and Oaks SC. 1992. *Emerging Infections: Microbial Threats to Health in the United States* (Washington, DC: National Academies Press, p. 64).
- ¹²⁰ Bäumlner AJ, Hargis BM, and Tsolis RM. 2000. Tracing the origins of *Salmonella* outbreaks. *Science* 287(5450):50-2.
- ¹²¹ Schroeder CM, Naugle AL, Schlosser WD, et al. 2005. Estimate of illnesses from *Salmonella* Enteritidis in eggs, United States, 2000. *Emerging Infectious Diseases* 11(1):113-5.
- ¹²² Morse EV and Duncan MA. 1974. Salmonellosis—an environmental health problem. *Journal of the American Veterinary Medical Association* 165(11):1015-9.
- ¹²³ Avens JS. 1987. Overview: *Salmonella*—what’s the problem? In: Colorado State University Poultry Symposium: Managing for Profit (Fort Collins, CO: Colorado State University, pp. 119-123).
- ¹²⁴ Bell DD. 2001. Cage management for layers. In: Bell DD and Weaver WD Jr (eds.), *Commercial Chicken Meat and Egg Production*, 5th Edition (Norwell, MA: Kluwer Academic Publishers, p 1009).
- ¹²⁵ United Egg Producers. 2010. United Egg Producers animal husbandry guidelines for U.S. egg laying flocks, 2010 edition. www.uepcertified.com/media/pdf/UEP-Animal-Welfare-Guidelines.pdf. Accessed March 15, 2010.
- ¹²⁶ Bell DD and Weaver WD (eds.). 2002. *Commercial Chicken Meat and Egg Production* (Norwell, Massachusetts: Kluwer Academic Publishers, p.1061).
- ¹²⁷ Greenfield J and Bankier JC. 1969. Isolation of *Salmonella* Arizona using enrichment media incubated at 35 and 43 C. *Avian Diseases* 13(4):864-71.
- ¹²⁸ World Health Organization. 2003. Technical paper: Main challenges in the control of zoonotic diseases in the Eastern Mediterranean Region. Agenda item 8(c). Regional Committee for the Eastern Mediterranean, fiftieth session. www.emro.who.int/rc50/documents/DOC7.doc. Accessed March 15, 2010.
- ¹²⁹ Hunter PR. 1992. Epizootics of *Salmonella* infection in poultry may be the result of modern selective breeding practices. *European Journal of Epidemiology* 8(6):851-5.
- ¹³⁰ Turnbull PCB. 1979. Food poisoning with special reference to *Salmonella*—its epidemiology, pathogenesis and control. *Clinics in Gastroenterology* 8(3):663-714.
- ¹³¹ Holt PS. 1993. Effect of induced molting on the susceptibility of White Leghorn hens to a *Salmonella* Enteritidis infection. *Avian Diseases* 37(2):412-7.
- ¹³² Clark GM, Kaufmann AF, Gangarosa EJ, and Thompson MA. 1973. Epidemiology of an international outbreak of *Salmonella* Agona. *The Lancet* 2(7827):490-3.
- ¹³³ Fritts CA, Kersey JH, and Waldroup PW. 2002. Utilization of spent hen meal in diets for laying hens. *International Journal of Poultry Science* 1(4):82-4.
- ¹³⁴ National Renderers Association. 2005. U.S. production, consumption and export of rendered products for 1998-2003.
- ¹³⁵ World Health Organization and the Office International des Epizooties. 1999. WHO consultation on public health and animal Transmissible Spongiform Encephalopathies: epidemiology, risk and research requirements (Geneva, Switzerland: December 1-3). www.who.int/csr/resources/publications/bse/WHO_CDS_CSRAPH_2000_2/en/. Accessed March 15, 2010.
- ¹³⁶ McChesney DG, Kaplan G, and Gardner P. 1995. FDA survey determines *Salmonella* contamination. *Feedstuffs* 67(7):20-3.
- ¹³⁷ Hirsch W and Sapiro-Hirsch R. 1958. The role of certain animal feeding stuffs especially bone meal, in the epidemiology of salmonellosis. *Harefuah* 54(3):59.
- ¹³⁸ Knox WA, Galbraith NS, Lewis MJ, Hickie GC, and Johnston HH. 1963. A milk-borne outbreak of food poisoning due to *Salmonella* Heidelberg. *The Journal of Hygiene* 61(2):175-85.
- ¹³⁹ Pennington JH, Brooksbank NH, Poole PM, and Seymour F. 1968. *Salmonella* Virchow in a chicken-packing station and associated rearing units. *British Medical Journal* 4(5634):804-6.

-
- ¹⁴⁰ Turnbull PCB. 1979. Food poisoning with special reference to *Salmonella*—its epidemiology, pathogenesis and control. *Clinics in Gastroenterology* 8(3):663-714.
- ¹⁴¹ Crump JA, Griffin PM, and Angulo FJ. 2002. Bacterial contamination of animal feed and its relationship to human foodborne illness. *Clinical Infectious Diseases* 35(7):859-65.
- ¹⁴² Hopp P, Wahlström H, and Hirn J. 1999. A common *Salmonella* control programme in Finland, Norway and Sweden. *Acta Veterinaria Scandinavica* 91(Suppl):45-9.
- ¹⁴³ Jacobson M and DeWaal CS. 1999. Testimony before the Senate Committee on Government Affairs on “Egg safety: are there cracks in the federal food safety system” (Washington, DC: July 1). www.cspinet.org/foodsafety/egg_safety.html. Accessed March 15, 2010.
- ¹⁴⁴ Drexler M. 2002. *Secret Agents: The Menace of Emerging Infections* (Washington, DC: Joseph Henry Press).
- ¹⁴⁵ Nestle M. 2003. *Safe Food: Bacteria, Biotechnology, and Bioterrorism* (Berkeley and Los Angeles, CA: University of California Press, p. 27).
- ¹⁴⁶ Fox N. 1997. *Spoiled: The Dangerous Truth about a Food Chain Gone Haywire* (New York, NY: Basic Books).
- ¹⁴⁷ Lewis C. 1998. *Safety last: the politics of E. coli and other food-borne killers*. Center for Public Integrity, February 26.
- ¹⁴⁸ Mead PS, Slutsker L, Dietz V, et al. 1999. Food-related illness and death in the United States. *Emerging Infectious Diseases* 5(5):607-25.
- ¹⁴⁹ Cheeke PR. 1999. *Contemporary Issues in Animal Agriculture*, 2nd Edition (Danville, IL: Interstate Publishers, Inc.).
- ¹⁵⁰ Pew Commission on Industrial Farm Animal Production. 2008. Putting meat on the table: industrial farm animal production in America. Executive summary, p. 13. www.ncifap.org/images/PCIFAPSmry.pdf. Accessed March 15, 2010.
- ¹⁵¹ Pew Commission on Industrial Farm Animal Production. 2008. Expert panel highlights serious public health threats from industrial animal agriculture. Press release issued April 11. www.pewtrusts.org/news_room_detail.aspx?id=37968. Accessed March 15, 2010.
- ¹⁵² Pew Commission on Industrial Farm Animal Production. 2008. Putting meat on the table: industrial farm animal production in America, p. 38. www.ncifap.org/images/PCIFAPFin.pdf. Accessed Accessed March 15, 2010.
- ¹⁵³ Pew Commission on Industrial Farm Animal Production. 2008. Putting meat on the table: industrial farm animal production in America. Executive summary, p. 13. www.ncifap.org/images/PCIFAPSmry.pdf. Accessed March 15, 2010.
- ¹⁵⁴ Pew Commission on Industrial Farm Animal Production. 2008. Pew Commission says industrial scale farm animal production poses “unacceptable” risks to public health, environment. Press release issued April 29.
- ¹⁵⁵ Cal. Health & Safety Code § 25990 *et seq.* (West 2009).
- ¹⁵⁶ MICH. COMP. LAWS § 287.746 (2010).
- ¹⁵⁷ California A.B. 1437 chaptered July 6, 2010. www.leginfo.ca.gov/cgi-bin/postquery?bill_number=ab_1437&sess=0910&house=A. Accessed 2 August 2010.

The Humane Society of the United States is the nation’s largest animal protection organization—backed by 11 million Americans, or one of every 28. For more than a half-century, The HSUS has been fighting for the protection of all animals through advocacy, education, and hands-on programs. Celebrating animals and confronting cruelty. On the Web at humanesociety.org.